

# **$p$ -ADIC INTERPOLATION FUNCTION RELATED TO MULTIPLE GENERALIZED GENOCCHI NUMBERS**

**SERKAN ARACI, MEHMET ACIKGOZ, AND ERDOĞAN ŞEN**

**ABSTRACT.** In the present paper, we deal with multiple generalized Genocchi numbers and polynomials. Also, we introduce analytic interpolating function for the multiple generalized Genocchi numbers attached to  $\chi$  at negative integers in complex plane and we define the multiple Genocchi  $p$ -adic  $L$ -function. Finally, we derive the value of the partial derivative of our multiple  $p$ -adic  $l$ -function at  $s = 0$ .

**2010 MATHEMATICS SUBJECT CLASSIFICATION.** 11S80, 11B68.

**KEYWORDS AND PHRASES.** Multiple generalized Genocchi numbers and polynomials, Euler-Gamma function,  $p$ -adic interpolation function, multiple generalized zeta function.

## **1. PRELIMINARIES**

The works of generalized Bernoulli, Euler and Genocchi numbers and polynomials and their combinatorial relations have received much attention [6], [7]-[27], [30], [31], [32], [33], [34]. Generalized Bernoulli polynomials, generalized Euler polynomials and generalized Genocchi numbers and polynomials are the signs of very strong relationship between elementary number theory, complex analytic number theory, Homotopy theory (stable Homotopy groups of spheres), differential topology (differential structures on spheres), theory of modular forms (Eisenstein series),  $p$ -adic analytic numbers theory ( $p$ -adic  $L$ -functions), quantum physics(quantum Groups).

$p$ -adic numbers also were invented by German Mathematician Kurt Hensel around the end of the nineteenth century. In spite of their being already one hundred years old, these numbers are still today enveloped in an aura of mystery within the scientific community. The  $p$ -adic integral was used in mathematical physics, for instance, the functional equation of the  $q$ -zeta function,  $q$ -stirling numbers and  $q$ -Mahler theory of integration with respect to the ring  $\mathbb{Z}_p$  together with Iwasawa's  $p$ -adic  $L$  functions.

Also the  $p$ -adic interpolation functions of the Bernoulli and Euler polynomials have been treated by Tsumura [35] and Young [36]. T. Kim [7]-[23] also studied on  $p$ -adic interpolation functions of these numbers and polynomials. In [37], Carlitz originally constructed  $q$ -Bernoulli numbers and polynomials. These numbers and polynomials are studied by many authors (see cf. [8]-[29], [41]). In the last decade, a surprising number of papers appeared proposing new generalizations of the Bernoulli, Euler and Genocchi polynomials to real and complex variables.

In [7]-[24], Kim studied some families of multiple Bernoulli, Euler and Genocchi numbers and polynomials. By using the fermionic  $p$ -adic invariant integral on  $\mathbb{Z}_p$ , he constructed  $p$ -adic Bernoulli,  $p$ -adic Euler and  $p$ -adic Genocchi numbers and polynomials of higher order.

In this paper, by using Kim's method in [20], we derive several properties for the multiple generalized Genocchi numbers attached to  $\chi$ .

As is well-known, Genocchi numbers are defined in the complex plane by the following exponential function

$$(1.1) \quad C(t) = \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad |t| < \pi.$$

It follows from the description that  $G_0 = 0$ ,  $G_1 = 1$ ,  $G_2 = -1$ ,  $G_3 = 0$ ,  $G_4 = 1$ ,  $G_5 = 0, \dots$ , and  $G_{2k+1} = 0$  for  $k = 1, 2, 3, \dots$ .

The Genocchi polynomials are also given by the rule:

$$C(t, x) = e^{tG(x)} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} = \frac{2t}{e^t + 1} e^{xt},$$

with the usual convention of replacing  $G^n(x) := G_n(x)$  (see [8], [9] and [11]).

Let  $w \in \mathbb{N}$ . Then the multiple Genocchi polynomials of order  $w$  are given by [24]

$$(1.2) \quad C^{(w)}(t, x) = \left( \frac{2t}{e^t + 1} \right)^w e^{xt} = \sum_{n=0}^{\infty} G_n^{(w)}(x) \frac{t^n}{n!}, \quad |t| < \pi.$$

Taking  $x = 0$  in (1.2), then we have  $G_n^{(w)}(0) := G_n^{(w)}$  are called the multiple Genocchi numbers of order  $w$ .

For  $f \in \mathbb{N}$  with  $f \equiv 1 \pmod{2}$ , we assume that  $\chi$  is a primitive Dirichlet's character with conductor  $f$ . It is known in [28] that the Genocchi numbers associated with  $\chi$ ,  $G_{n,\chi}$ , was introduced by the following expression

$$(1.3) \quad C_\chi(t) = 2t \sum_{\xi=1}^f \frac{\chi(\xi) (-1)^\xi e^{\xi t}}{e^{ft} + 1} = \sum_{n=0}^{\infty} G_{n,\chi} \frac{t^n}{n!}, \quad |t| < \frac{\pi}{f}.$$

In this paper, we contemplate the definition of the generating function of the multiple generalized Genocchi numbers attached to  $\chi$  in the complex plane. From this definition, we introduce an analytic interpolating function for the multiple generalized Genocchi numbers attached to  $\chi$ . Finally, we investigate behaviour of analytic interpolating function at  $s = 0$ .

## 2. ON AN ANALYTIC FUNCTION IN CONNECTION WITH THE MULTIPLE GENERALIZED GENOCCHI NUMBERS

In this part, we introduce the multiple generalized Genocchi numbers attached to  $\chi$  defined by

$$(2.1) \quad \begin{aligned} C_\chi^{(w)}(t) &= \sum_{n=0}^{\infty} G_{n,\chi}^{(w)} \frac{t^n}{n!} \\ &= (2t)^w \sum_{a_1, \dots, a_w=1}^f \frac{(-1)^{a_1+\dots+a_w} \chi(a_1+\dots+a_w) e^{t(a_1+\dots+a_w)}}{(e^{ft} + 1)^w}. \end{aligned}$$

On account of (1.2) and (2.1), we easily derive the following

$$(2.2) \quad G_{n,\chi}^{(w)} = \frac{f^n}{f^w} \sum_{a_1, \dots, a_w=1}^f (-1)^{a_1+\dots+a_w} \chi(a_1+\dots+a_w) G_n^{(w)} \left( \frac{a_1+\dots+a_w}{f} \right).$$

For  $s \in \mathbb{C}$ , we have

$$(2.3) \quad \begin{aligned} & \frac{1}{\Gamma(s)} \int_0^\infty t^{s-w-1} \left\{ (-1)^w C^{(w)}(-t, x) \right\} dt \\ &= 2^w \sum_{n_1, \dots, n_w \geq 0} \frac{(-1)^{n_1 + \dots + n_w}}{(x + n_1 + \dots + n_w)^s}, \quad x \neq 0, -1, -2, \dots \end{aligned}$$

where  $\Gamma(s)$  is Euler-Gamma function, which is defined by the rule

$$\Gamma(s) = \int_0^\infty t^{s-1} e^{-t} dt.$$

Thanks to (2.3), we give the multiple Genocchi-zeta function as follows: for  $s \in \mathbb{C}$  and  $x \neq 0, -1, -2, \dots$ ,

$$(2.4) \quad \zeta_G^{(w)}(s, x) = 2^w \sum_{n_1, \dots, n_w \geq 0} \frac{(-1)^{n_1 + \dots + n_w}}{(x + n_1 + \dots + n_w)^s}.$$

By (1.2) and (2.3), we see that

$$\zeta_G^{(w)}(-n, x) = \frac{G_{n+w}^{(w)}(x)}{\binom{n+w}{w} w!} \text{ for } n \in \mathbb{N}.$$

By utilizing from complex integral and (2.1), we obtain the following equation: for  $s \in \mathbb{C}$ ,

$$(2.5) \quad \begin{aligned} & \frac{1}{\Gamma(s)} \int_0^\infty t^{s-w-1} \left\{ (-1)^w C_\chi^{(w)}(-t) \right\} dt \\ &= 2^w \sum_{\substack{n_1, \dots, n_w = 0 \\ n_1 + \dots + n_w \neq 0}} \frac{\chi(a_1 + \dots + a_w) (-1)^{n_1 + \dots + n_w}}{(n_1 + \dots + n_w)^s}, \end{aligned}$$

where  $\chi$  is the primitive Dirichlet's character with conductor

$$f \in \mathbb{N} \text{ and } f \equiv 1 \pmod{2}.$$

Because of (2.5), we give the definition Dirichlet's type of multiple Genocchi  $L$ -function in complex plane as follows:

$$(2.6) \quad L^{(w)}(s | \chi) = 2^w \sum_{\substack{n_1, \dots, n_w = 0 \\ n_1 + \dots + n_w \neq 0}}^\infty \frac{\chi(a_1 + \dots + a_w) (-1)^{n_1 + \dots + n_w}}{(n_1 + \dots + n_w)^s}.$$

Via the (2.1) and (2.6), we derive the following theorem:

**Theorem 1.** *For any  $n \in \mathbb{N}$ , then we have*

$$(2.7) \quad L^{(w)}(-n | \chi) = \frac{G_{n+w, \chi}^{(w)}(x)}{\binom{n+w}{w} w!}.$$

Let  $s$  be a complex variable, and let  $a$  and  $b$  be integer with  $0 < a < F$  and  $F \equiv 1 \pmod{2}$ .

Thus, we can consider the partial zeta function  $S^{(w)}(s; a_1, \dots, a_w \mid F)$  as follows:

$$\begin{aligned}
 (2.8) \quad & S^{(w)}(s; a_1, \dots, a_w \mid F) \\
 &= 2^w \sum_{\substack{m_1, \dots, m_w > 0 \\ m_i \equiv a_i \pmod{F}}} \frac{(-1)^{m_1 + \dots + m_w}}{(m_1 + \dots + m_w)^s} \\
 &= (-1)^{a_1 + \dots + a_w} F^{-s} \zeta_G^{(w)} \left( s, \frac{a_1 + \dots + a_w}{F} \right).
 \end{aligned}$$

**Theorem 2.** *The following identity holds:*

$$S^{(w)}(s; a_1, \dots, a_w \mid F) = (-1)^{a_1 + \dots + a_w} F^{-s} \zeta_G^{(w)} \left( s, \frac{a_1 + \dots + a_w}{F} \right).$$

Then Dirichlet's type of multiple  $L$ -function can be expressed as the sum: for  $s \in \mathbb{C}$

$$(2.9) \quad L^{(w)}(s \mid \chi) = \sum_{a_1, \dots, a_w = 1}^F \chi(a_1 + \dots + a_w) S^{(w)}(s; a_1, \dots, a_w \mid F).$$

Substituting  $s = w - n$  into (2.8), we readily derive the following: for  $w, n \in \mathbb{N}$

$$\begin{aligned}
 (2.10) \quad & \binom{n}{w} w! S^{(w)}(w - n; a_1, \dots, a_w \mid F) \\
 &= (-1)^{a_1 + \dots + a_w} F^{n-w} G_n^{(w)} \left( \frac{a_1 + \dots + a_w}{F} \right).
 \end{aligned}$$

By (1.2), it is simple to indicate the following

$$(2.11) \quad G_n^{(w)}(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} G_k^{(w)} = \sum_{k=0}^n \binom{n}{k} x^k G_{n-k}^{(w)}.$$

Thanks to (2.8), (2.10) and (2.11), we develop the following theorem:

**Theorem 3.** *The following identity*

$$\begin{aligned}
 (2.12) \quad & w! \binom{-s}{w} S^{(w)}(s + w; a_1, \dots, a_w \mid F) \\
 &= (-1)^{a_1 + \dots + a_w} F^{-w} (a_1 + \dots + a_w)^{-s} \sum_{k \geq 0} \binom{-s}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}
 \end{aligned}$$

is true.

From (2.9), (2.10) and (2.12), we have the following corollary:

**Corollary 1.** *The following holds true:*

$$\begin{aligned}
 (2.13) \quad & w! \binom{-s}{w} L^{(w)}(s + w \mid \chi) \\
 &= \sum_{a_1, \dots, a_w = 1}^F \chi(a_1 + \dots + a_w) (-1)^{a_1 + \dots + a_w} F^{-w} (a_1 + \dots + a_w)^{-s} \\
 &\quad \times \sum_{k=0}^{\infty} \binom{-s}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}.
 \end{aligned}$$

The values of  $L^{(w)}(s | \chi)$  at negative integers are algebraic, hence may be regarded as lying in an extension of  $\mathbb{Q}_p$ . We therefore look for a  $p$ -adic function which agrees with  $L^{(w)}(s | \chi)$  at the negative integers in the next section.

### 3. CONCLUSION

In this final section, we consider  $p$ -adic interpolation function of the multiple generalized Genocchi  $L$ -function, which interpolate Dirichlet's type of multiple Genocchi numbers at negative integers. Firstly, Washington constructed  $p$ -adic  $l$ -function which interpolates generalized classical Bernoulli numbers.

Here, we use some the following notations, which will be useful in remainder of paper.

Let  $\omega$  denote the *Teichmüller* character by the conductor  $f_\omega = p$ . For an arbitrary character  $\chi$ , we set  $\chi_n = \chi\omega^{-n}$ ,  $n \in \mathbb{Z}$ , in the sense of product of characters.

Let

$$\langle a \rangle = \omega^{-1}(a) a = \frac{a}{\omega(a)}.$$

Thus, we note that  $\langle a \rangle \equiv 1 \pmod{p\mathbb{Z}_p}$ . Let

$$A_j(x) = \sum_{n=0}^{\infty} a_{n,j} x^n, \quad a_{n,j} \in \mathbb{C}_p, \quad j = 0, 1, 2, \dots$$

be a sequence of power series, each convergent on a fixed subset

$$T = \left\{ s \in \mathbb{C}_p \mid |s|_p < p^{-\frac{2-p}{p-1}} \right\},$$

of  $\mathbb{C}_p$  such that

- (1)  $a_{n,j} \rightarrow a_{n,0}$  as  $j \rightarrow \infty$  for any  $n$ ;
- (2) for each  $s \in T$  and  $\epsilon > 0$ , there exists an  $n_0 = n_0(s, \epsilon)$  such that

$$\left| \sum_{n \geq n_0} a_{n,j} s^n \right|_p < \epsilon \text{ for } \forall j.$$

So,

$$\lim_{j \rightarrow \infty} A_j(s) = A_0(s), \text{ for all } s \in T.$$

This was firstly introduced by Washington [43] to indicate that each functions  $\omega^{-s}(a) a^s$  and

$$\sum_{k=0}^{\infty} \binom{s}{k} \left( \frac{F}{a} \right)^k B_k,$$

where  $F$  is multiple of  $p$  and  $f$  and  $B_k$  is the  $k$ -th Bernoulli numbers, is analytic on  $T$  (for more information, see [43]).

We assume that  $\chi$  is a primitive Dirichlet's character with conductor  $f \in \mathbb{N}$  with  $f \equiv 1 \pmod{2}$ . Then we contemplate the multiple Genocchi  $p$ -adic  $L$ -function,  $L_p^{(w)}(s | \chi)$ , which interpolates the multiple generalized Genocchi numbers attached to  $\chi$  at negative integers.

For  $f \in \mathbb{N}$  with  $f \equiv 1 \pmod{2}$ , let us assume that  $F$  is a positive integral multiple of  $p$  and  $f = f_\chi$ . We now give the definition of mutiple Genocchi  $p$ -adic  $L$ -function

as follows:

$$\begin{aligned}
 (3.1) \quad & w! \binom{-s}{w} L_p^{(w)}(s+w \mid \chi) \\
 &= \frac{1}{F^w} \sum_{a_1, \dots, a_w=1}^F \chi(a_1 + \dots + a_w) (-1)^{a_1 + \dots + a_w} \langle a_1 + \dots + a_w \rangle^{-s} \\
 &\quad \times \sum_{k=0}^{\infty} \binom{-s}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}.
 \end{aligned}$$

Due to (3.1), we want to note that  $L_p^{(w)}(s+w \mid \chi)$  is an analytic function on  $s \in T$ .

For  $n \in \mathbb{N}$ , we have

$$(3.2) \quad G_{n, \chi_n}^{(w)} = \frac{F^n}{F^w} \sum_{a_1, \dots, a_w=1}^F (-1)^{a_1 + \dots + a_w} \chi_n(a_1 + \dots + a_w) G_n^{(w)} \left( \frac{a_1 + \dots + a_w}{F} \right).$$

If  $\chi_n(p) \neq 0$ , then  $(p, f_{\chi_n}) = 1$ , and so the ratio  $\frac{F}{p}$  is a multiple of  $f_{\chi_n}$ .

Let

$$\vartheta = \left\{ \frac{a_1 + \dots + a_w}{p} \mid a_1 + \dots + a_w \equiv 0 \pmod{p} \text{ for some } a_i \in \mathbb{Z} \text{ with } 0 \leq a_i \leq F \right\}.$$

Therefore we can write the following

$$\begin{aligned}
 (3.3) \quad & \frac{F^n}{F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ p \mid a_1 + \dots + a_w}}^F (-1)^{a_1 + \dots + a_w} \chi_n(a_1 + \dots + a_w) G_n^{(w)} \left( \frac{a_1 + \dots + a_w}{F} \right) \\
 &= p^{n-w} \frac{\left(\frac{F}{p}\right)^n}{\left(\frac{F}{p}\right)^w} \chi_n(p) \sum_{\substack{a_1, \dots, a_w=1 \\ \lambda \in \vartheta}}^{\frac{F}{p}} (-1)^\lambda \chi_n(\lambda) G_n^{(w)} \left( \frac{\lambda}{F/p} \right).
 \end{aligned}$$

By (3.3), we define the different multiple generalized Genocchi numbers attached to  $\chi$  as follows:

$$(3.4) \quad G_{n, \chi_n}^{*(w)} = \frac{\left(\frac{F}{p}\right)^n}{\left(\frac{F}{p}\right)^w} \sum_{\substack{a_1, \dots, a_w=1 \\ \lambda \in \vartheta}}^{\frac{F}{p}} (-1)^\lambda \chi_n(\lambda) G_n^{(w)} \left( \frac{\lambda}{F/p} \right).$$

On account of (3.2), (3.3) and (3.4), we attain the following

$$(3.5) \quad G_{n, \chi_n}^{(w)} - p^{n-w} \chi_n(p) G_{n, \chi_n}^{*(w)} = \frac{F^n}{F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ p \nmid a_1 + \dots + a_w}}^F (-1)^{a_1 + \dots + a_w} \chi_n(a_1 + \dots + a_w) G_n^{(w)} \left( \frac{a_1 + \dots + a_w}{F} \right).$$

By the definition of the multiple Genocchi polynomials of order  $w$ , we write the following

$$(3.6) \quad G_n^{(w)} \left( \frac{a_1 + \dots + a_w}{F} \right) = F^{-n} (a_1 + \dots + a_w)^n \sum_{k=0}^n \binom{n}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}.$$

By (3.5) and (3.6), we have

$$\begin{aligned}
 (3.7) \quad & G_{n, \chi_n}^{(w)} - p^{n-w} \chi_n(p) G_{n, \chi_n}^{*(w)} \\
 &= \frac{1}{F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ p \nmid a_1 + \dots + a_w}}^F (-1)^{a_1 + \dots + a_w} \chi_n(a_1 + \dots + a_w) (a_1 + \dots + a_w)^n \\
 &\quad \times \sum_{k=0}^n \binom{n}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}
 \end{aligned}$$

By (3.1) and (3.7), we readily see that

$$\begin{aligned}
 & w! \binom{n}{w} L_p^{(w)}(w - n \mid \chi) \\
 &= \frac{1}{F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ p \nmid a_1 + \dots + a_w}}^F (-1)^{a_1 + \dots + a_w} \chi_n(a_1 + \dots + a_w) (a_1 + \dots + a_w)^n \\
 (3.8) \quad & \times \sum_{k=0}^n \binom{n}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)} \\
 &= G_{n, \chi_n}^{(w)} - p^{n-w} \chi_n(p) G_{n, \chi_n}^{*(w)}
 \end{aligned}$$

Consequently, we arrive at the following theorem.

**Theorem 4.** *The following nice identity holds true:*

$$\begin{aligned}
 & w! \binom{-s}{w} L_p^{(w)}(s + w \mid \chi) \\
 &= \frac{1}{F^w} \sum_{a_1, \dots, a_w=1}^F \chi(a_1 + \dots + a_w) (-1)^{a_1 + \dots + a_w} \langle a_1 + \dots + a_w \rangle^{-s} \\
 &\quad \times \sum_{k=0}^{\infty} \binom{-s}{k} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)}.
 \end{aligned}$$

Thus  $L_p^{(w)}(s + w \mid \chi)$  is an analytic function on  $T$ . Additionally, for each  $n \in \mathbb{N}$ , we procure the following:

$$w! \binom{n}{w} L_p^{(w)}(w - n \mid \chi) = G_{n, \chi_n}^{(w)} - p^{n-w} \chi_n(p) G_{n, \chi_n}^{*(w)}.$$

Using Taylor expansion at  $s = 0$ , we have

$$(3.9) \quad \binom{-s}{k} = \frac{(-1)^k}{k} s + \dots \text{ if } k \geq 1.$$

Differentiating on both sides in (3.1), with respect to  $s$  at  $s = 0$ , we obtain the following corollary.

**Theorem 5.** *Let  $F$  be a positive integral multiple of  $p$  and  $f$ . Then we have*

$$\begin{aligned} & \frac{\partial}{\partial s} \left( \binom{-s}{w} L_p^{(w)}(s+w \mid \chi) \right) \Big|_{s=0} = \frac{(-1)^w}{w} L_p^{(w)}(w \mid \chi) \\ &= \frac{1}{w!F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ (a_1+\dots+a_w, p)=1}}^F \chi(a_1 + \dots + a_w) (-1)^{a_1+\dots+a_w} (1 - \log_p(a_1 + \dots + a_w)) \\ &+ \frac{1}{w!F^w} \sum_{\substack{a_1, \dots, a_w=1 \\ (a_1+\dots+a_w, p)=1}}^F \chi(a_1 + \dots + a_w) (-1)^{a_1+\dots+a_w} \sum_{m=1}^{\infty} \frac{(-1)^m}{m} \left( \frac{F}{a_1 + \dots + a_w} \right)^k G_k^{(w)} \end{aligned}$$

where  $\log_p x$  is the  $p$ -adic logarithm.

## REFERENCES

- [1] Araci, S., Erdal, D., and Seo, J.-J., A study on the fermionic  $p$ -adic  $q$ -integral representation on  $\mathbb{Z}_p$  associated with weighted  $q$ -Bernstein and  $q$ -Genocchi polynomials, *Abstract and Applied Analysis*, Volume **2011**, Article ID 649248, 10 pages.
- [2] Araci, S., Seo, J.-J., and Erdal, D., New construction weighted  $(h, q)$ -Genocchi numbers and polynomials related to Zeta type functions, *Discrete Dynamics in Nature and Society*, Volume **2011**, Article ID 487490, 7 pages.
- [3] Araci, S., Acikgoz, M., and Seo, J. J., Explicit formulas involving  $q$ -Euler numbers and polynomials, *Abstract and Applied Analysis*, Volume **2012**, Article ID 298531, 11 pages.
- [4] Araci, S., Acikgoz, M., and Jolany, H., On  $p$ -adic interpolating function associated with modified Dirichlet's type of twisted  $q$ -euler numbers and polynomials with weight alpha, *submitted*.
- [5] Araci, S., Acikgoz, M., and Park, K. H., A note on the  $q$ -analogue of Kim's  $p$ -adic log gamma type functions associated with  $q$ -extension of Genocchi and Euler numbers with weight alpha, *Bulletin of the Korean Mathematical Society* (In press).
- [6] Araci, S., Acikgoz, M., Park, K. H., and Jolany, H., On the unification of two families of multiple twisted type polynomials by using  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  at  $q = -1$ , *Bulletin of the Malaysian Mathematical Sciences and Society* (In press).
- [7] Kim, T., On a  $p$ -adic interpolation function for the  $q$ -extension of the generalized Bernoulli polynomials and its derivative, *Discrete Mathematics*, 309 (2009) 1593–1602.
- [8] Kim, T., On the  $q$ -extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.* 326 (2007) 1458–1465.
- [9] Kim, T., On the multiple  $q$ -Genocchi and Euler numbers, *Russian J. Math. Phys.* 15 (4) (2008) 481–486.
- [10] Kim, T.,  $p$ -adic  $q$ -integrals associated with Changhee-Barnes'  $q$ -Bernoulli polynomials, *Integral Transforms Spec. Funct.* 15 (2004) 415–420.
- [11] Kim, T., A Note on the  $q$ -Genocchi Numbers and Polynomials, *Journal of Inequalities and Applications* Article ID 71452, 8 pages.
- [12] Kim, T.,  $q$ -Volkenborn integration, *Russ. J. Math. phys.* 9 (2002), 288–299.
- [13] Kim, T., An invariant  $p$ -adic  $q$ -integrals on  $\mathbb{Z}_p$ , *Applied Mathematics Letters*, vol. 21 (2008), pp. 105–108.
- [14] Kim, T.,  $q$ -Euler numbers and polynomials associated with  $p$ -adic  $q$ -integrals, *J. Nonlinear Math. Phys.*, 14 (2007), no. 1, 15–27.
- [15] Kim, T., New approach to  $q$ -Euler polynomials of higher order, *Russ. J. Math. Phys.*, 17 (2010), no. 2, 218–225.
- [16] Kim, T., Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ , *Russ. J. Math. Phys.*, 16 (2009), no.4, 484–491.
- [17] Kim, T. and Rim, S.-H., On the twisted  $q$ -Euler numbers and polynomials associated with basic  $q$ - $l$ -functions, *Journal of Mathematical Analysis and Applications*, vol. 336 (2007), no. 1, pp. 738–744.
- [18] Kim, T., On  $p$ -adic  $q$ - $l$ -functions and sums of powers, *J. Math. Anal. Appl.* (2006).



- [19] Kim, T., On the analogs of Euler numbers and polynomials associated with  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  at  $q = -1$ , *J. Math. Anal. Appl.* 331 (2) (2007) 779-792.
- [20] Kim, T., On  $p$ -adic interpolating function for  $q$ -Euler numbers and its derivatives, *J. Math. Anal. Appl.* 339 (1) (2008) 598-608.
- [21] Kim, T.,  $q$ -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients, *Russ. J. Math. Phys.* 15 (1) (2008) 51-57.
- [22] Kim, T., Euler numbers and polynomials associated with zeta functions, *Abstr. Appl. Anal.* (2008) Art. ID 581582, 11 pp.
- [23] Kim, T., Analytic continuation of multiple  $q$ -zeta functions and their values at negative integers, *Russ. J. Math. Phys.* 11 (1) (2004) 71-76.
- [24] Ryoo, C. S., Kim, T., Choi, J., and Lee, B., On the generalized  $q$ -Genocchi numbers and polynomials of higher-order, *Advances in Difference Equations*, vol. 2011, Article ID 424809, 8 pages, 2011.
- [25] Kang, J.Y., Lee, H. Y., Jung, N.S., Ryoo, C.S., On the Barnes' type related to multiple Genocchi polynomials on  $\mathbb{Z}_p$ , *Abstract and Applied Analysis*, Volume 2012, Article ID 679495, 9 pages.
- [26] Kang, J. Y., Some properties of multiple generalized  $q$ -Genocchi polynomials with weight  $\alpha$  and weak weight  $\beta$ , *International Journal of Mathematics and Mathematical Sciences*, Volume 2012, Article ID 179385, 13 pages.
- [27] Jang, L-C., and Kim, T.,  $q$ -Genocchi numbers and polynomials associated with fermionic  $p$ -adic invariant integrals on  $\mathbb{Z}_p$ , *Abstr. Appl. Anal.* (2008) Art. ID 232187, 8 pp.
- [28] Jang, L.-C., Hwang, K.-W., and Kim, Y.-H., A note on  $(h, q)$ -Genocchi polynomials and numbers of higher order, *Advances in Difference Equations*, vol. 2010, Article ID 309480, 6 pages, 2010.
- [29] Park, Kyoung Ho., On Interpolation Functions of the Generalized Twisted  $(h, q)$ -Euler Polynomials, *Journal of Inequalities and Applications*, Volume 2009, Article ID 946569, 17 pages
- [30] G. D. Liu, Generating functions and generalized Euler numbers, *Proc. Japan Acad. Ser. A Math. Sci.*, 84 (2008), 29-34.
- [31] C. Frappier, Representation formulas for entire functions of exponential type and Generalized Bernoulli polynomials, *J. Austral. Math. Soc. Ser.*, 64 (1998), No. 3, 307-316.
- [32] Jolany, H., and Darafsheh, M. R., Some another remarks on the generalization of Bernoulli and Euler numbers, *Scientia Magna*, Vol. 5 (2009), No. 3, 118-129.
- [33] Jolany, H., Alikelaye, R. Eizadi., and Mohamad, S-H., Some results on the generalization of Bernoulli, Euler and Genocchi polynomials, *Acta Universitatis Apulensis*, No. 27 (2011), 299-306.
- [34] Vandiver, H-S., On Generalizations of the Numbers of Bernoulli and Euler, Vol. 23, No. 10 (1937), 555-559.
- [35] Tsumura, H., On a  $p$ -adic interpolation of the generalized Euler numbers and it applications, *Tokyo J. Math.* 10 (2) (1987) 281-293.
- [36] Young, P., Congruences for Bernoulli, Euler, and Stirling numbers, *J. Number Theory* 78 (1999) 204-227.
- [37] Carlitz, L.,  $q$ -Bernoulli and Eulerian numbers, *Trans. Amer. Math. Soc.* 76 (1954) 332-370.
- [38] Acikgoz, M., and Simsek, Y., On multiple interpolation function of the Nörlund-type  $q$ -Euler polynomials, *Abst. Appl. Anal.* 2009 (2009), Article ID 382574, 14 pages.
- [39] Ozden, H., and Simsek, Y., A new extension of  $q$ -Euler numbers and polynomials related to their interpolation functions, *Appl. Math. Lett.* 21 (2008) 934-939.
- [40] Shiratani, K., and Yamamoto, S., On a  $p$ -adic interpolation function for the Euler numbers and its derivatives, *Mem. Fac. Sci. Kyushu Univ. Ser. A* 39 (1) (1985) 113-125.
- [41] Srivastava, H-M., Kim, T., and Simsek, Y.,  $q$ -Bernoulli numbers and polynomials associated with multiple  $q$ -zeta functions and basic  $L$ -series, *Russ. J. Math. Phys.* 12 (2005) 241-268.
- [42] Srivastava, H-M., and Pintér, Á., Remarks on some relationships between the Bernoulli and Euler polynomials, *Appl. Math. Lett.* 17 (2004) 375-380.
- [43] Washington, L-C., Introduction to cyclotomic fields, Second edition, *Springer-Verlag*, 1997.
- [44] G. J. Fox, A method of Washington applied to the derivation of a two-variable  $p$ -adic  $L$ -function, *Pacific J. Math.* 209 (2003), 31-40.
- [45] Diamond, J., On the values of  $p$ -adic  $L$ -functions at positive integers, *Acta Arith.* 35 (1979), 223-237.

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,  
27310 Gaziantep, TURKEY

*E-mail address:* mtsrkn@hotmail.com; mtsrkn@gmail.com; saraci88@yahoo.com.tr

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,  
27310 Gaziantep, TURKEY

*E-mail address:* acikgoz@gantep.edu.tr

Department of Mathematics, Faculty of Science and Letters, Namik Kemal Uni-  
versity, 59030 Tekirdağ, TURKEY

*E-mail address:* erdogan.math@gmail.com